

JBMF-22-69993

Exposed roots of *Salvadora oleoides* in Aravalli range addresses freeways constraint: ecological and management perspective

Maneesh S Bhandari^{1*}, Rajendra K Meena¹, Aman Dabral¹, Jaspal S Chauhan² and Shailesh Pandey³

¹Department Genetics and Tree Improvement, Forest Research Institute, Uttarakhand, India

²Department of Aquatic Biodiversity, Hemwati Nandan Bahuguna Central University, Uttarakhand, India

³Department of Forest Protection, Forest Research Institute, Uttarakhand, India

***Corresponding author:** Maneesh S Bhandari, Department Genetics and Tree Improvement, Forest Research Institute, Uttarakhand, India, Tel: +911352224383; E-mail: maneesh31803@gmail.com

Received date: 22 July, 2022, Manuscript No. JBMF-22-69993;

Editor assigned date: 25 July, 2022, PreQC No. JBMF-22-69993 (PQ);

Reviewed date: 08 August, 2022, QC No. JBMF-22-69993;

Revised date: 03 October, 2022, Manuscript No. JBMF-22-69993 (R);

Published date: 10 October, 2022, DOI: 10.4172/2327-4417.1000025.

Supplementary Material



(a)



(b)



(c)



(d)



(e)



(f)



(g)



(h)



(i)



(j)



(k)



(l)



(m)

Figure 1: List of events showing extraction and digging of soil, impacts on *S. oleoides* and its arid biodiversity (a) – (m): a) Dhandlan digged community land (Jhajjar); b) Digged land (Matanhail, Jhajjar); c) marks of tractor trolley; d) Mustard field near Jawa (Bhiwani); e) *S. oleoides* exposed root; f) Ruminants of roots; g) Logged *S.oleoides*; h) Stone crusher machine; i) Crushed sand for truck loading; j) Domesticated animals lying in *S. oleoides* community forest land; k) Exposed barren land showed *Pavo cristatus*; l) Parrot near Rudrol (Bhiwani); m) Avian diversit.